

2010

Ozsváth-Szabó and Rasmussen Invariants of Cable Knots

Cornelia A. Van Cott

University of San Francisco, cvancott@usfca.edu

Follow this and additional works at: <http://repository.usfca.edu/math>

 Part of the [Geometry and Topology Commons](#)

Recommended Citation

Cornelia A Van Cott. Ozsváth–Szabó and Rasmussen invariants of cable knots. *Algebraic & Geometric Topology*, 10 (2010) 825–836
DOI: 10.2140/agt.2010.10.825

This Article is brought to you for free and open access by the College of Arts and Sciences at USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. It has been accepted for inclusion in Mathematics by an authorized administrator of USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. For more information, please contact repository@usfca.edu.

Ozsváth–Szabó and Rasmussen invariants of cable knots

CORNELIA A VAN COTT

We study the behavior of the Ozsváth–Szabó and Rasmussen knot concordance invariants τ and s on $K_{m,n}$, the (m,n) –cable of a knot K where m and n are relatively prime. We show that for every knot K and for any fixed positive integer m , both of the invariants evaluated on $K_{m,n}$ differ from their value on the torus knot $T_{m,n}$ by fixed constants for all but finitely many $n > 0$. Combining this result together with Hedden’s extensive work on the behavior of τ on $(m, mr + 1)$ –cables yields bounds on the value of τ on any (m,n) –cable of K . In addition, several of Hedden’s obstructions for cables bounding complex curves are extended.

57M25

1 Introduction

The (m,n) –cable of a knot K , denoted $K_{m,n}$, is the satellite knot with companion K and pattern $T_{m,n}$, the (m,n) –torus knot. The behavior of many classical concordance invariants has been shown to be rather predictable on cable knots. For example, it is a classical result (see Lickorish [6]) that the Alexander polynomial of a cable knot is

$$\Delta_{K_{m,n}}(t) = \Delta_K(t^m)\Delta_{T_{m,n}}(t).$$

Shinohara [17] found a formula for the signature of a cable knot, and Litherland [7] extended the result, finding the value of Tristram–Levine signatures on a cable knot:

$$\sigma_\omega(K_{m,n}) = \sigma_\omega^m(K) + \sigma_\omega(T_{m,n}).$$

Milnor signatures and Casson–Gordon invariants of cables (see Litherland [8] and Kearton [5], respectively, for details) also yield nice formulas.

The purpose of this note is to investigate two relatively new concordance invariants – the Ozsváth–Szabó invariant τ and the Rasmussen invariant s – and their behavior on cable knots. The discussion here will use only the formal properties that the two invariants have in common.

Both τ and s were introduced in connection with developments in the theory of knot homologies: τ is defined in terms of knot Floer homology (see Ozsváth and Szabó [10] and Rasmussen [15]) and the Rasmussen invariant s is defined in terms of Khovanov homology (see Rasmussen [14]). These two invariants have enabled important progress

in the field of knot theory, providing new proofs for Milnor's conjecture [10; 14] and examples of Alexander polynomial one knots which are not smoothly slice (see Livingston [9]).

No work has been done to compute the Rasmussen invariant for cables, but the behavior of the Ozsváth–Szabó concordance invariant τ under $(m, mr + 1)$ -cabling has been investigated by Hedden [2; 3]. Through careful investigation of the relationship between the filtered chain homotopy types of $\mathcal{F}(K_{m, mr+1}, i)$ and $\mathcal{F}(K, i)$, he obtained the following main result:

Theorem 1 [3] *Let $K \subset S^3$ be a nontrivial knot. Then the following inequality holds for all r :*

$$m\tau(K) + \frac{(mr)(m-1)}{2} \leq \tau(K_{m, mr+1}) \leq m\tau(K) + \frac{(mr)(m-1)}{2} + m - 1.$$

In the special case when K satisfies $\tau(K) = g(K)$, we have the equality

$$\tau(K_{m, mr+1}) = m\tau(K) + \frac{(mr)(m-1)}{2},$$

whereas when $\tau(K) = -g(K)$, we have

$$\tau(K_{m, mr+1}) = m\tau(K) + \frac{(mr)(m-1)}{2} + m - 1.$$

When appropriately normalized, τ and s share several formal properties and agree on many families of knots, though in general they have been shown to be distinct invariants (see Hedden and Ording [4]). Stated in reference to τ , the essential formal properties are as follows (see Ozsváth and Szabó [10]):

- (1) τ is a homomorphism from the smooth knot concordance group \mathcal{C} to \mathbb{Z} .
- (2) $|\tau(K)| \leq g_4(K)$, where $g_4(K)$ denotes the 4-genus of K .
- (3) $\tau(T_{m,n}) = (m-1)(n-1)/2$, where $T_{m,n}$ denotes the (m, n) -torus knot with $m, n \geq 1$.

It can be shown that $s/2$ also satisfies these three properties [14]. In addition, both τ and s are insensitive to a change in orientation [11; 14]. Our main results will only depend on these formal properties, and hence apply to both invariants. To proceed concisely, let ν denote any concordance invariant satisfying the above properties.

Fixing $m > 0$, we would like to study the value of ν on $K_{m,n}$ as a function of n , where n ranges over the integers relatively prime to m . (Notice that $K_{m,n} = -K_{-m,-n}$, and so the restriction $m > 0$ does not limit our results.) From our observations about other concordance invariants, we expect that the behavior of $\nu(K_{m,n})$ as a function of n is somehow related to the behavior of $\nu(T_{m,n})$. This, in fact, is true. As a function of n ,

$v(T_{m,n})$ is linear of slope $(m - 1)/2$ for $n > 0$. We will see that the function $v(K_{m,n})$ is close to being linear with the same slope. Specifically, we subtract from v a linear function to construct the following function:

$$h(n) = v(K_{m,n}) - \frac{(m - 1)}{2}n,$$

where n is an integer relatively prime to m . We have the following theorem:

Theorem 2 *The function $h(n)$ is a nonincreasing $\frac{1}{2} \cdot \mathbb{Z}$ -valued function which is bounded below. In particular, we have*

$$-(m - 1) \leq h(n) - h(r) \leq 0$$

for all $n > r$, where both n and r are relatively prime to m .

From this result it follows that for all n large enough, h is constant. Hence for n large enough, $v(K_{m,n})$ differs from $v(T_{m,n})$ by a fixed constant. That is, for every knot K there exist integers N and c such that $v(K_{m,n}) = v(T_{m,n}) + c$ for all $n > N$, where n is relatively prime to m . Additionally, a similar statement with corresponding constant c' holds for all $n < N'$ for some N' .

Theorem 2 is sharp in the sense that there are knots K with associated functions h which achieve the bounds given in the theorem. For example, when K is slice, $h(n) = (m - 1)/2$ for all $n < 0$ and $h(n) = -(m - 1)/2$ for all $n > 0$. Here the drop in functional value from $n = -1$ to $n = 1$ is maximal: $h(1) - h(-1) = -(m - 1)$. On the other hand, we will see that when $v = \tau$ and $\tau(K) = g_3(K)$, the function h is constant.

Using **Theorem 2**, we can take several results which apply only to $(m, mr + 1)$ -cables and extend their scope to include *all* cables. For example, the bounds on the value of τ on $(m, mr + 1)$ -cables described in **Theorem 1** extend to all cables as follows.

Corollary 3 *Let $K \subset S^3$ be a nontrivial knot. Then the following inequality holds for all n relatively prime to m :*

$$m\tau(K) + \frac{(m - 1)(n - 1)}{2} \leq \tau(K_{m,n}) \leq m\tau(K) + \frac{(m - 1)(n + 1)}{2}.$$

When K satisfies $\tau(K) = g(K)$, we have

$$\tau(K_{m,n}) = m\tau(K) + \frac{(m - 1)(n - 1)}{2},$$

whereas when $\tau(K) = -g(K)$, we have

$$\tau(K_{m,n}) = m\tau(K) + \frac{(m - 1)(n + 1)}{2}.$$

Observe that the results in [Corollary 3](#) could probably also have been obtained by using the definition of τ and studying the filtered chain homotopy type of $\mathcal{F}(K_{m,n})$ for n relatively prime to m . However, the proof here avoids this and uses only the analysis of $\mathcal{F}(K_{m,mr+1})$ in [\[3\]](#) together with [Theorem 2](#) to obtain the result for all cables.

The second half of [Corollary 3](#) motivates studying knots K for which $\tau(K) = g(K)$. Hedden summarized many results about such knots and their $(m, mr + 1)$ -cables in [\[3\]](#). Now combining that discussion with [Corollary 3](#) from above, we can extend several of his results to a more general setting. Let \mathcal{P} denote the class of all knots satisfying the equality $\tau(K) = g(K)$. An immediate consequence of [Corollary 3](#) is the following.

Corollary 4 *Let K be a nontrivial knot in S^3 , and let n be relatively prime to m .*

- (1) *If $K \in \mathcal{P}$, then $K_{m,n} \in \mathcal{P}$ if and only if $n > 0$.*
- (2) *If $K \notin \mathcal{P}$, then $K_{m,n} \notin \mathcal{P}$.*

As discussed in [\[3\]](#), \mathcal{P} contains several classes of knots. We mention two such classes here:

- Any knot K which bounds a properly embedded complex curve, $V_f \subset B^4$, with $g(V_f) = g(K)$. This set of knots includes, for example, positive knots (that is, knots which admit diagrams with only positive crossings). (See Hedden [\[1\]](#) and Livingston [\[9\]](#).)
- Any knot which admits a positive lens space (or L-space) surgery. (See Ozsváth and Szabó [\[12\]](#).)

From this, we have the following immediate applications extending the work of [\[3\]](#).

Corollary 5 *If $K_{m,n}$ bounds a properly embedded complex curve $V_f \subset B^4$ satisfying $g(V_f) = g(K_{m,n})$, then $n > 0$ and $\tau(K) = g(K)$.*

Corollary 6 *Suppose that $K_{m,n}$ admits a positive lens space (or L-space) surgery. Then $n > 0$ and $\tau(K) = g(K)$.*

Corollary 7 *Suppose $K \notin \mathcal{P}$. Then $K_{m,n}$ is not a positive knot for any relatively prime pair of integers m, n .*

A final corollary concerns a more general class of knots – the class of \mathbb{C} -knots. A knot K is a \mathbb{C} -knot if K bounds a properly embedded complex curve $V_f \subset B^4$. From [\[1; 13; 16\]](#), we know that for such knots, $\tau(K) = g_4(K) \geq 0$. Coupling this result with [Corollary 3](#), we have the following corollary.

Corollary 8 Suppose that $K_{m,n}$ is a \mathbb{C} -knot. Then $n \geq -2m\tau(K)/(m-1) - 1$.

The primary significance of each of these corollaries is that they can be used as obstructions to cables having the discussed properties. Moreover, it is interesting that τ provides obstructions to such a wide array of geometric notions. For an excellent extended discussion of this, we refer the reader to [3].

This paper is organized as follows. Section 2 contains the proof of Theorem 2. Section 3 contains the proof of Corollary 3. Finally, in Section 4 we observe that the strategy for the proof of Theorem 2 extends to a broader setting in which, instead of cabling, we consider a braiding construction.

Acknowledgments I thank both Charles Livingston and Matthew Hedden for several helpful conversations.

2 Proof of Theorem 2

Let r, n be integers relatively prime to m with $n > r$. The general strategy here is to first find a cobordism between $K_{m,n} \# -K_{m,r}$ and a torus knot.

We begin with the knot $K_{m,n} \# -K_{m,r}$. Working through signs and orientations carefully, we find that

$$K_{m,n} \# -K_{m,r} = K_{m,n} \# (-K)_{m,-r}.$$

We will now do a series of band moves to the knot $K_{m,n} \# (-K)_{m,-r}$. A band move on any knot $K \subset S^3$ is accomplished as follows. Start with an embedding $b: I \times I \rightarrow S^3$ such that $b(I \times I) \cap K = b(I \times \{0, 1\})$ and such that b respects the orientation of K . Define $K_b = K - b(I \times \{0, 1\}) \cup b(\{0, 1\} \times I)$. The knot (or link) K_b is the result of doing a band move along b . Doing a band move to a knot simultaneously constructs a cobordism from the knot K to K_b . The genus of this cobordism can be computed explicitly. For example, in the special case that the result of performing a sequence of band moves is again a knot, one can show that the genus of the cobordism is half of the number of bands added.

Now there is a sequence of $m-1$ band moves on $K_{m,n} \# (-K)_{m,-r}$ which results in the knot (or link) $(K \# -K)_{m,n-r}$. See Figure 1 for an example. Since $K \# -K$ is cobordant to the unknot, $(K \# -K)_{m,n-r}$ is cobordant to the torus link $T_{m,n-r}$. Let k_+ denote the smallest positive integer such that $n-r+k_+$ is relatively prime to m . (If $n-r$ is already relatively prime to m , then set $k_+ = 0$.) By doing $k_+ \cdot (m-1)$ additional band moves to the torus link $T_{m,n-r}$, we obtain the torus knot $T_{m,n-r+k_+}$.




Figure 1: Beginning with the knot $K_{3,2} \# (-K)_{3,-1}$, we perform two band moves and obtain the knot $(K \# -K)_{3,1} \cdot K^m$. K^m denotes the mirror image of K .

(Figure 2). Altogether, the total number of band moves performed was $(k_+ + 1)(m - 1)$. Therefore, the knot $K_{m,n} \# -K_{m,r}$ is genus $(k_+ + 1)(m - 1)/2$ cobordant to the torus knot $T_{m,n-r+k_+}$. Hence we conclude that

$$g_4(K_{m,n} \# -K_{m,r} \# -T_{m,n-r+k_+}) \leq \frac{(k_+ + 1)(m - 1)}{2}.$$

Now since $|\nu(K)| \leq g_4(K)$, it follows that

$$|\nu(K_{m,n} \# -K_{m,r} \# -T_{m,n-r+k_+})| \leq \frac{(k_+ + 1)(m - 1)}{2}.$$




Figure 2: Beginning with the torus knot $T_{3,2}$, we perform two band moves and obtain $T_{3,3}$.

Simplifying the expression using the properties of ν , we obtain

$$\left| \nu(K_{m,n}) - \nu(K_{m,r}) - \frac{(m-1)(n-r+k_+-1)}{2} \right| \leq \frac{(k_++1)(m-1)}{2}.$$

At this point, recall the function $h(n)$ which we defined earlier. Using the definition of h , we can further simplify the inequality:

$$\left| h(n) - h(r) - \frac{(m-1)(k_+-1)}{2} \right| \leq \frac{(k_++1)(m-1)}{2}.$$

Hence,

$$(1) \quad -(m-1) \leq h(n) - h(r) \leq k_+(m-1).$$

Notice that if $k_+ = 0$, then we are done. If not, then we continue as follows.

Similar to before, let k_- denote the largest negative integer such that $n-r+k_-$ is relatively prime to m . By doing $|k_-| \cdot (m-1)$ band moves to $T_{m,n-r}$, we can obtain the torus knot $T_{m,n-r+k_-}$. Proceeding through the same steps as before, we obtain

$$(2) \quad (k_- - 1)(m-1) \leq h(n) - h(r) \leq 0.$$

Combining (1) and (2), we have

$$-(m-1) \leq h(n) - h(r) \leq 0$$

for all integers $n > r$ where both n and r are relatively prime to m .

3 Proof of Corollary 3

Combining Theorem 1 and Theorem 2 together, we obtain an easy proof that the bounds on the value of τ on $(m, mr+1)$ -cables described in Theorem 1 extend to all cables. We now restate and prove Corollary 3.

Corollary 3 *Let $K \subset S^3$ be a nontrivial knot. Then the following inequality holds for all n relatively prime to m :*

$$m\tau(K) + \frac{(m-1)(n-1)}{2} \leq \tau(K_{m,n}) \leq m\tau(K) + \frac{(m-1)(n+1)}{2}.$$

When K satisfies $\tau(K) = g(K)$, we have

$$\tau(K_{m,n}) = m\tau(K) + \frac{(m-1)(n-1)}{2},$$

whereas when $\tau(K) = -g(K)$, we have

$$\tau(K_{m,n}) = m\tau(K) + \frac{(m-1)(n+1)}{2}.$$

Proof The proof of this corollary is obtained by carefully combining the equalities and inequalities found in Theorem 1 and Theorem 2. We will demonstrate a portion of the proof, leaving the rest to the reader.

Let m and n be two relatively prime integers with $m > 0$. Let r be an integer such that $n > mr + 1$. Then by Theorem 2,

$$h(n) - h(mr + 1) \leq 0.$$

Using the definition of h and letting $v = \tau$, we obtain

$$\tau(K_{m,n}) \leq \tau(K_{m,mr+1}) - \frac{m-1}{2}(mr - n + 1).$$

Using the upper bound on $\tau(K_{m,mr+1})$ given by Theorem 1, we have

$$\tau(K_{m,n}) \leq m\tau(K) + \frac{(m-1)(n+1)}{2},$$

which is one side of the desired inequality.

To obtain the other side of the inequality, let r' be an integer such that $mr' + 1 > n$. Then by [Theorem 2](#),

$$h(mr' + 1) - h(n) \leq 0.$$

We leave to the reader the task of reducing this inequality (using methods exactly similar to above) to obtain the desired second half of the inequality in the corollary.

Now let K be a knot such that $\tau(K) = g(K)$. Suppose for contradiction that $\tau(K_{m,n}) \neq m\tau(K) + (m - 1)(n - 1)/2$. By the inequality discussed above, this implies that $\tau(K_{m,n}) > m\tau(K) + (m - 1)(n - 1)/2$. Again, let r be an integer such that $n > mr + 1$. Then we have

$$\begin{aligned} h(n) - h(mr + 1) &= \tau(K_{m,n}) - \frac{(m - 1)}{2}n - \tau(K_{m,mr+1}) + \frac{m - 1}{2}(mr + 1) \\ &= \tau(K_{m,n}) - \frac{(m - 1)}{2}n - m\tau(K) + \frac{(m - 1)}{2} \\ &> m\tau(K) + \frac{(m - 1)(n - 1)}{2} - \frac{(m - 1)}{2}n - m\tau(K) + \frac{(m - 1)}{2} \\ &= 0. \end{aligned}$$

This contradicts [Theorem 2](#). Therefore, $\tau(K_{m,n}) = m\tau(K) + (m - 1)(n - 1)/2$ for all n relatively prime to m . A similar argument settles the case when K is a knot such that $\tau(K) = -g(K)$. □

4 Further analysis

The process of cabling a knot can be reinterpreted as a special case of the following more general procedure. Let β be an element of the braid group B_m such that the closure of the braid $\hat{\beta}$ is a knot. There is a natural solid torus V which contains the closed braid $\hat{\beta}$. Remove a neighborhood of a knot K in S^3 and glue in the solid torus V by a homeomorphism which maps longitude to longitude and meridian to meridian. We denote the resulting knot by K_β . Notice that if we take the braid $\beta \in B_m$ to be $(\sigma_{m-1}\sigma_{m-2} \cdots \sigma_1)^n$ (where σ_i denotes the i -th standard generator of the braid group), then the resulting knot K_β is the (m, n) -cable $K_{m,n}$.

For any braid $\beta \in B_m$, let β_r denote the braid consisting of β with r full twists adjoined to the end of the braid. Specifically, $\beta_r = \beta(\sigma_{m-1}\sigma_{m-2} \cdots \sigma_1)^{mr}$. The value of ν on K_{β_r} as a function of r turns out to have controlled behavior similar to that of cabling. Define the function

$$g(r) = \nu(K_{\beta_r}) - \frac{(m - 1)}{2}mr,$$

where $\beta \in B_m$ is a braid whose closure is a knot and r is an integer. Then we have the following theorem about the behavior of the function g .

Theorem 9 *The function $g(r)$ is a nonincreasing integer valued function which is bounded below. In particular,*

$$-(m-1) \leq g(r) - g(s) \leq 0$$

for all $r > s$.

From this theorem, it follows that the function g is eventually constant. This allows us to describe quite clearly a relationship among the values of τ (and s) on an entirely new set of knots. Fixing a knot K and a braid $\beta \in B_m$ such that $\hat{\beta}$ is a knot, [Theorem 9](#) implies that for all large r ,

$$v(K_{\beta_{r+1}}) = v(K_{\beta_r}) + \frac{m(m-1)}{2}.$$

where v can be taken to be either τ or s . Note that if we take K in the above construction to be the unknot, then the theorem relates the values of v on knots with braid representatives which differ by full twists.

We turn now to the proof of [Theorem 9](#).

Proof As with the proof of [Theorem 2](#), the first goal here is to find a cobordism between $K_{\beta_r} \# -K_{\beta_s}$ and a torus knot. Notice that $-K_{\beta_s} = (-K)_{(\beta^{-1})_{-s}}$. Therefore,

$$K_{\beta_r} \# -K_{\beta_s} = K_{\beta_r} \# (-K)_{(\beta^{-1})_{-s}}.$$

By doing $m-1$ band moves to the latter knot, we obtain the knot $(K \# -K)_{(\beta\beta^{-1})_{r-s}}$. Since $K \# -K$ is cobordant to the unknot and $\beta\beta^{-1}$ is the trivial m -strand braid, this new knot is cobordant to the torus link $T_{m,m(r-s)}$. Again, by doing $(m-1)$ band moves to the torus link $T_{m,m(r-s)}$, we obtain the torus knot $T_{m,m(r-s)+1}$. A total of $2(m-1)$ band moves have been performed. Therefore, the knot $K_{\beta_r} \# -K_{\beta_s}$ is genus $(m-1)$ cobordant to the torus knot $T_{m,m(r-s)+1}$. Hence

$$g_4(K_{\beta_r} \# -K_{\beta_s} \# -T_{m,m(r-s)+1}) \leq m-1.$$

Now since $|v(K)| \leq g_4(K)$, it follows that

$$|v(K_{\beta_r} \# -K_{\beta_s} \# -T_{m,m(r-s)+1})| \leq m-1,$$

which simplifies to

$$\left| v(K_{\beta_r}) - v(K_{\beta_s}) - \frac{(m-1)m(r-s)}{2} \right| \leq m-1.$$

We now recall the function $g(r)$ which we defined earlier. Using the definition of g , we can further simplify the inequality and obtain

$$(3) \quad -(m-1) \leq g(r) - g(s) \leq m-1.$$

This gives us only half of the desired inequality. To obtain the remaining half, go back to the torus link $T_{m,m(r-s)}$ which we obtained from $K_{\beta_r} \# -K_{\beta_s}$ by a cobordism which added $m-1$ bands. Instead of adding $m-1$ additional bands to obtain the torus knot $T_{m,m(r-s)+1}$, add $m-1$ bands to obtain the torus knot $T_{m,m(r-s)-1}$. Proceeding through the same steps as before, we obtain

$$(4) \quad -2(m-1) \leq g(r) - g(s) \leq 0.$$

Combining (3) and (4), we have

$$-(m-1) \leq g(r) - g(s) \leq 0$$

for all integers $r > s$, as desired. \square

References

- [1] **M Hedden**, *Notions of positivity and the Ozsváth–Szabó concordance invariant* [arXiv:math/0509499](#)
- [2] **M Hedden**, *On knot Floer homology and cabling*, *Algebr. Geom. Topol.* 5 (2005) 1197–1222 [MR2171808](#)
- [3] **M Hedden**, *On knot Floer homology and cabling. II*, *Int. Math. Res. Not.* (2009) 2248–2274 [MR2511910](#)
- [4] **M Hedden, P Ording**, *The Ozsváth–Szabó and Rasmussen concordance invariants are not equal*, *Amer. J. Math.* 130 (2008) 441–453 [MR2405163](#)
- [5] **C Kearton**, *The Milnor signatures of compound knots*, *Proc. Amer. Math. Soc.* 76 (1979) 157–160 [MR534409](#)
- [6] **WBR Lickorish**, *An introduction to knot theory*, *Graduate Texts in Math.* 175, Springer, New York (1997) [MR1472978](#)
- [7] **RA Litherland**, *Signatures of iterated torus knots*, from: “Topology of low-dimensional manifolds (Proc. Second Sussex Conf., Chelwood Gate, 1977)”, (R A Fenn, editor), *Lecture Notes in Math.* 722, Springer, Berlin (1979) 71–84 [MR547456](#)
- [8] **RA Litherland**, *Cobordism of satellite knots*, from: “Four-manifold theory (Durham, N.H., 1982)”, (C Gordon, R Kirby, editors), *Contemp. Math.* 35, Amer. Math. Soc. (1984) 327–362 [MR780587](#)
- [9] **C Livingston**, *Computations of the Ozsváth–Szabó knot concordance invariant*, *Geom. Topol.* 8 (2004) 735–742 [MR2057779](#)

- [10] **P Ozsváth, Z Szabó**, *Knot Floer homology and the four-ball genus*, *Geom. Topol.* 7 (2003) 615–639 [MR2026543](#)
- [11] **P Ozsváth, Z Szabó**, *Holomorphic disks and knot invariants*, *Adv. Math.* 186 (2004) 58–116 [MR2065507](#)
- [12] **P Ozsváth, Z Szabó**, *On knot Floer homology and lens space surgeries*, *Topology* 44 (2005) 1281–1300 [MR2168576](#)
- [13] **O Plamenevskaya**, *Bounds for the Thurston–Bennequin number from Floer homology*, *Algebr. Geom. Topol.* 4 (2004) 399–406 [MR2077671](#)
- [14] **J A Rasmussen**, *Khovanov homology and the slice genus* [arXiv:math/0402131](#)
- [15] **J A Rasmussen**, *Floer homology and knot complements*, PhD thesis, Harvard University (2003)
- [16] **L Rudolph**, *Algebraic functions and closed braids*, *Topology* 22 (1983) 191–202 [MR683760](#)
- [17] **Y Shinohara**, *On the signature of knots and links*, *Trans. Amer. Math. Soc.* 156 (1971) 273–285 [MR0275415](#)

Department of Mathematics, University of San Francisco
San Francisco, California, 94117

cvancott@usfca.edu

Received: 28 December 2009