

2017

Use of Flow Cytometry and Stable Isotope Analysis to Determine Phytoplankton Uptake of Wastewater Derived Ammonium in a Nutrient-rich River

Calla Schmidt

University of San Francisco, cischmidt@usfca.edu

Tamara E. C. Kraus

Megan B. Young

Carol Kendall

Follow this and additional works at: <https://repository.usfca.edu/envs>

Recommended Citation

Schmidt, Calla; Kraus, Tamara E. C.; Young, Megan B.; and Kendall, Carol, "Use of Flow Cytometry and Stable Isotope Analysis to Determine Phytoplankton Uptake of Wastewater Derived Ammonium in a Nutrient-rich River" (2017). *Environmental Science*. 38. <https://repository.usfca.edu/envs/38>

This Article is brought to you for free and open access by the College of Arts and Sciences at USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. It has been accepted for inclusion in Environmental Science by an authorized administrator of USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. For more information, please contact repository@usfca.edu.

Use of flow cytometry and stable isotope analysis to determine phytoplankton uptake of wastewater derived ammonium in a nutrient-rich river

Calla M. Schmidt¹, Tamara E. C. Kraus², Megan B. Young³, Carol Kendall³

¹University of San Francisco, 2130 Fulton St, San Francisco, CA, 94117, USA

²USGS California Water Science Center, 6000 J Street, Placer Hall, Sacramento CA 95819, USA

³USGS National Research Program, Menlo Park, CA, USA

Correspondence to: Calla M Schmidt (cis Schmidt@usfca.edu)

Abstract. Anthropogenic alteration of the form and concentration of nitrogen (N) in aquatic ecosystems is widespread. Understanding availability and uptake of different N sources at the base of aquatic food webs is critical to establishment of effective nutrient management programs. Stable isotopes of N (¹⁴N, ¹⁵N) are often used to trace the sources of N fuelling aquatic primary production, but effective use of this approach requires obtaining a reliable isotopic ratio for phytoplankton. In this study, we tested the use of flow cytometry to isolate phytoplankton from bulk particulate organic matter (POM) in a portion of the Sacramento River, California, during river-scale nutrient manipulation experiments that involved halting wastewater discharges high in ammonium (NH₄⁺). Field samples were collected using a Lagrangian approach, allowing us to measure changes in phytoplankton N source in the presence and absence of wastewater derived NH₄⁺. Comparison of δ¹⁵N-POM and δ¹⁵N-phytoplankton (δ¹⁵N-PHY) revealed that their δ¹⁵N values followed broadly similar trends. However, after 3 days of downstream travel in the presence of wastewater treatment plant (WWTP) effluent, δ¹⁵N-POM and δ¹⁵N-PHY in the Sacramento River differed by as much as 7‰. Using a stable isotope mixing model approach, we estimated that in the presence of effluent between 40 and 90% of phytoplankton-N was derived from NH₄⁺ after 3 days of downstream transport. An apparent gradual increase over time in the proportion of NH₄⁺ in the phytoplankton N pool suggests that either very low phytoplankton growth rates resulted in an N turnover time that exceeded the travel time sampled during this study, or a portion of the phytoplankton community continued to access nitrate even in the presence of elevated NH₄⁺ concentrations.

