

B.M.A.T. - Banner Mobility Assessment Tool for Nurses

Test	Task	Response	Fail = Choose Most Appropriate Equipment/Device(s)	Pass
Assessment Level 1 Assessment of: -Cognition -Trunk strength -Seated balance	<p>Sit and Shake: From a semi-reclined position, ask patient to sit upright and rotate* to a seated position at the side of the bed; <i>may use the bedrail.</i></p> <p>Note patient's ability to maintain bedside position.</p> <p>Ask patient to reach out and grab your hand and shake making sure patient reaches across his/her midline.</p> <p>Note: Consider your patients cognitive ability, including orientation and CAM assessment if applicable.</p>	<p>Sit: Patient is able to follow commands, has some trunk strength; caregivers may be able to try weight-bearing if patient is able to maintain seated balance greater than two minutes (without caregiver assistance).</p> <p>Shake: Patient has significant upper body strength, awareness of body in space, and grasp strength.</p>	<p>MOBILITY LEVEL 1</p> <ul style="list-style-type: none"> - Use total lift with sling and/or repositioning sheet and/or straps. - Use lateral transfer devices such as roll board, friction reducing (slide sheets/tube), or air assisted device. <p>NOTE: <i>If patient has 'strict bed rest' or bilateral 'non-weight bearing' restrictions, do not proceed with the assessment; patient is MOBILITY LEVEL 1.</i></p>	Passed Assessment Level 1 = Proceed with Assessment Level 2.
Assessment Level 2 Assessment of : -Lower extremity strength -Stability	<p>Stretch and Point: With patient in seated position at the side of the bed, have patient place both feet on the floor (or stool) with knees no higher than hips.</p> <p>Ask patient to stretch one leg and straighten the knee, then bend the ankle/flex and point the toes. If appropriate, repeat with the other leg.</p>	<p>Patient exhibits lower extremity stability, strength and control.</p> <p>May test only one leg and proceed accordingly (e.g., stroke patient, patient with ankle in cast).</p>	<p>MOBILITY LEVEL 2</p> <ul style="list-style-type: none"> - Use total lift for patient unable to weight-bear on at least one leg. - Use sit-to-stand lift for patient who can weight-bear on at least one leg. 	Passed Assessment Level 2 = Proceed with Assessment Level 3.
Assessment Level 3 Assessment of: -Lower extremity strength for standing	<p>Stand: Ask patient to elevate off the bed or chair (seated to standing) using an assistive device (cane, bedrail).</p> <p>Patient should be able to raise buttocks off bed and hold for a count of five. May repeat once.</p> <p>Note: Consider your patients cognitive ability, including orientation and CAM assessment if applicable.</p>	<p>Patient exhibits upper and lower extremity stability and strength.</p> <p>May test with weight-bearing on only one leg and proceed accordingly (e.g., stroke patient, patient with ankle in cast).</p> <p>If any assistive device (cane, walker, crutches) is needed, patient is Mobility Level 3.</p>	<p>MOBILITY LEVEL 3</p> <ul style="list-style-type: none"> - Use non-powered raising/stand aid; <i>default to powered sit-to-stand lift if no stand aid available.</i> - Use total lift with ambulation accessories. - Use assistive device (cane, walker, crutches). <p>NOTE: Patient passes Assessment Level 3 but requires assistive device to ambulate or cognitive assessment indicates poor safety awareness; <i>patient is MOBILITY LEVEL 3.</i></p>	Passed Assessment Level 3 AND no assistive device needed = Proceed with Assessment Level 4. <p style="text-align: center;">Consult with Physical Therapist when needed and appropriate.</p>
Assessment Level 4 Assessment of: -Standing balance -Gait	<p>Walk: Ask patient to march in place at bedside. Then ask patient to advance step and return each foot.</p> <p>Patient should display stability while performing tasks. Assess for stability and safety awareness.</p>	<p>Patient exhibits steady gait and good balance while marching, and when stepping forwards and backwards.</p> <p>Patient can maneuver necessary turns for in-room mobility.</p> <p>Patient exhibits safety awareness.</p>	<p>MOBILITY LEVEL 3</p> <p>If patient shows signs of unsteady gait or fails Assessment Level 4, refer back to MOBILITY LEVEL 3; patient is MOBILITY LEVEL 3.</p>	<p>MOBILITY LEVEL 4</p> <p>MODIFIED INDEPENDENCE</p> <p>Passed = No assistance needed to ambulate; use your best clinical judgment to determine need for supervision during ambulation.</p>

Always default to the safest lifting/transfer method (e.g., total lift) if there is any doubt in the patient's ability to perform the task.