


Fall 12-16-2016

# Pharmacovigilance in Post-Marketing: Risk Assessment and Reporting Standards

Tiffany Jee

University of San Francisco, jee.tiffany@gmail.com

Follow this and additional works at: <https://repository.usfca.edu/capstone>

 Part of the [Other Public Health Commons](#), and the [Pharmacy Administration, Policy and Regulation Commons](#)

---

## Recommended Citation

Jee, Tiffany, "Pharmacovigilance in Post-Marketing: Risk Assessment and Reporting Standards" (2016). *Master's Projects and Capstones*. 557.

<https://repository.usfca.edu/capstone/557>

This Project/Capstone is brought to you for free and open access by the Theses, Dissertations, Capstones and Projects at USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. It has been accepted for inclusion in Master's Projects and Capstones by an authorized administrator of USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. For more information, please contact [repository@usfca.edu](mailto:repository@usfca.edu).

**Pharmacovigilance in Post-Marketing: Risk Assessment and Reporting Standards**

**Tiffany Jee**

**MPH Candidate 2016**

**University of San Francisco**

## Abstract

While there has been new targeted drug therapies released into the market over the recent years, pulmonary arterial hypertension (PAH) remains to be a rapidly progressive disease with poor prognosis. Over time the arteries stiffen and tighten subsequently leaving the heart to pump harder to try and provide enough blood to the body. The extra stress of the increased pumping leads to weakening and enlargement of the heart. The endothelin targeted therapy, Letairis (ambrisentan), is an oral tablet proven to improve quality of life in patients suffering from PAH, WHO Group I – functional class II and III by helping the arteries loosen and relax so that blood can flow steadily to the rest of the body. Although Letairis improves the quality of life for patients, there is a known serious risk of teratogenicity for females of reproductive potential. The risk of teratogenicity is so prominent that the Food and Drug Administration (FDA) required Letairis be placed on the risk evaluation and mitigation strategy program (REMS) to continually assess the risks and benefits of the product during post-marketing. In order to prevent and reduce harm in patients, mechanisms for evaluation and monitoring are vital to drug and patient safety.

*Keywords: pharmacovigilance, adverse event, pulmonary arterial hypertension, risk evaluation and mitigation strategy, ambrisentan, letairis*

## **I. Introduction**

Pulmonary arterial hypertension (PAH) is a rare progressive disease that affects people of all ages, races, and ethnic backgrounds and is caused by the narrowing and/or tightening of the pulmonary arteries, which connect the right side of the heart to the lungs. In many cases, as PAH develops in patients, blood flow through the pulmonary arteries is restricted and the right side of the heart becomes enlarged due to the increased strain of blood being pumped through the lungs and into the rest of the body. PAH is considered a chronic disease which currently lacks a cure. The World Health Organization (WHO) has classified pulmonary hypertension (PH) into 5 groups based on its causes. Group 1 is divided into 2 subgroups: PAH occurring with a known cause (Associated PAH) and PAH occurring without a known cause (Idiopathic PAH). PH Groups 2 to 5 are referred to as “Secondary PH” because the PH is most likely due to a pre-existing condition such as left heart disease, hypoxemia, chronic thrombotic pulmonary hypertension (CTEPH), or multifactorial mechanisms (Humbert, Sitbon, & Simonneau, 2004). Many who suffer from PAH are misdiagnosed or diagnosed (typically at age  $50 \pm 15$  years) after the disease has aggressively progressed to an advanced stage leaving the patient with few options for treatment (Stamm, Risbano, & Mathier, 2011).

During the 1980's, those living with PAH, especially those with idiopathic PAH, were finding that the disease would rapidly progress and lead to heart failure or death and patients would see a 2.8-year median survival rate after initial diagnosis (Humbert et al, 2004). Some of the first few prospective PAH studies during this time period also showed that the actuarial survival rate after diagnosis was 68-77% at 1-year, 40-56% at 3-years, and 22-38% at 5-years respectively (Humbert et al, 2004). The prevalence of PAH has increased over the years to 15-50 cases per million, substantially effecting those of certain high risk groups the most (PAH-

info.com, 2016). Associated and Secondary PAH have a prevalence of 0.5% in patients infected with human immunodeficiency virus (HIV), 2-3.75% in patients with sickle cell disease, and 7-12% in patients with systemic sclerosis (PAH-info.com, 2016). Idiopathic PAH (IPAH) has an incidence rate of 1-2 cases per million however, it is 2 to 4 times more likely to occur in women than in men and is thought to account for at least 40% of PAH cases overall (PAH-info.com, 2016). This IPAH statistic is also supported by the Pulmonary Hypertension Association Registry (PHAR) which lists 39% of those registered since 2015 are characterized as being diagnosed with IPAH (PHAssociation.org, 2016).

*\*The full version of the original paper is on file with the Director of the MPH Program.*

### Peer Reviewed Articles

1. Figueiras, A., Tato, F., Fontaiñas, J., & Gestal-Otero, J. J. (1999). Influence of physicians' attitudes on reporting adverse drug events: a case-control study. *Medical care*, 37 (8), 809-814.
2. Galiè, N., Barberà, J. A., Frost, A. E., Ghofrani, H. A., Hoeper, M. M., McLaughlin, V. V., ... & Oudiz, R. J. (2015). Initial Use of Ambrisentan Plus Tadalafil in Pulmonary Arterial Hypertension. *New England Journal of Medicine*, 373 (9), 834-844.
3. Hassoun, P. M., Zamanian, R. T., Damico, R., Lechtzin, N., Khair, R., Kolb, T. M., ... & Sato, T. (2015). Ambrisentan and Tadalafil Up-Front Combination Therapy in Scleroderma-Associated Pulmonary Arterial Hypertension. *American Journal of Respiratory and Critical Care Medicine*, 192(9), 1102-1110.

### References

1. American Heart Association (2015). Pulmonary Hypertension. Retrieved November 18, 2016 from [http://www.heart.org/HEARTORG/Conditions/CongenitalHeartDefects/TheImpactofCongenitalHeartDefects/Pulmonary-Hypertension\\_UCM\\_307044\\_Article.jsp-.WDz3aqIrLdQ](http://www.heart.org/HEARTORG/Conditions/CongenitalHeartDefects/TheImpactofCongenitalHeartDefects/Pulmonary-Hypertension_UCM_307044_Article.jsp-.WDz3aqIrLdQ)
2. Chick, A. and Magnuson, D. (2015). Risk Evaluation and Mitigation Strategy [PowerPoint Presentation].
3. Gilead (2016). Gilead Sciences Inc.. Retrieved November 21, 2016 from <http://www.gilead.com/about>
4. Horton, L. (2007). Food and Drug Administration. *BMJ: British Medical Journal*, 334(7584), 55-56. Retrieved from <http://0-www.jstor.org.ignacio.usfca.edu/stable/20506073>

5. Humbert, M., Sitbon, O., & Simonneau, G. (2004). Treatment of Pulmonary Arterial Hypertension. *New England Journal of Medicine*, 351(14), 1425-1436.
6. National Heart, Lung, and Blood Institute (2011). Pulmonary Hypertension. Retrieved November 18, 2016 from <http://www.nhlbi.nih.gov/health/health-topics/topics/pah/>
7. PAH-info (2016). Understanding PH. Retrieved November 21, 2016 from <http://www.pah-info.com/Home>
8. Pulmonary Hypertension Association (2016). Pulmonary Hypertension Association Registry: Preliminary Data. Retrieved on November 18, 2016 from <http://www.phassociation.org/PHAR/PreliminaryData>
9. Stamm, J. A., Risbano, M. G., & Mathier, M. A. (2011). Overview of Current Therapeutic Approaches for Pulmonary Hypertension. *Pulmonary circulation*, 1(2), 138-159.
10. U.S. Food & Drug Administration (2016). Regulatory Information. Retrieved November 22, 2016 from <http://www.fda.gov/RegulatoryInformation/Legislation/SignificantAmendmentstotheFDCAct/FoodandDrugAdministrationAmendmentsActof2007/default.htm>
11. World Health Organization. (2004). Pharmacovigilance: ensuring the safe use of medicines. *WHO/EDM/2004.8*. Geneva: World Health Organization.